

Package ‘MetabolicSyndrome’

January 20, 2025

Type Package

Title Diagnosis of Metabolic Syndrome

Version 0.1.3

Description

The modified Adult Treatment Panel -III guidelines (ATP-III) proposed by American Heart Association (AHA) and National Heart, Lung and Blood Institute (NHLBI) are used widely for the clinical diagnosis of Metabolic Syndrome. The AHA-NHLBI criteria advise using parameters such as waist circumference (WC), systolic blood pressure (SBP), diastolic blood pressure (DBP), fasting plasma glucose (FPG), triglycerides (TG) and high-density lipoprotein cholesterol (HDL-C) for diagnosis of metabolic syndrome. Each parameter has to be interpreted based on the proposed cut-offs, making the diagnosis slightly complex and error-prone. This package is developed by incorporating the modified ATP-III guidelines, and it will aid in the easy and quick diagnosis of metabolic syndrome in busy healthcare settings and also for research purposes. The modified ATP-III-AHA-NHLBI criteria for the diagnosis is described by Grundy et al ., (2005) <[doi:10.1161/CIRCULATIONAHA.105.169404](https://doi.org/10.1161/CIRCULATIONAHA.105.169404)>.

License GPL-3

Encoding UTF-8

LazyData true

RoxygenNote 7.2.3

URL <https://github.com/jagadishramasamy/metsynd>

BugReports <https://github.com/jagadishramasamy/metsynd/issues>

Suggests knitr, rmarkdown

Imports dplyr

Depends R (>= 2.10)

NeedsCompilation no

Author Jagadish Ramasamy [aut, cre] (<<https://orcid.org/0000-0003-4725-3227>>)

Maintainer Jagadish Ramasamy <iamjagankmr@gmail.com>

Repository CRAN

Date/Publication 2023-08-18 18:12:32 UTC

Contents

MetabolicSyndrome	2
x	3
Index	4

MetabolicSyndrome	<i>Diagnosis of Metabolic Syndrome</i>
-------------------	----------------------------------------

Description

The modified Adult Treatment Panel -III guidelines (ATP-III) proposed by American Heart Association (AHA) and National Heart, Lung and Blood Institute (NHLBI) are used widely for the clinical diagnosis of Metabolic Syndrome. The AHA-NHLBI criteria advise using parameters such as waist circumference (WC), systolic blood pressure (SBP), diastolic blood pressure (DBP), fasting plasma glucose (FPG), triglycerides (TG) and high-density lipoprotein cholesterol (HDL) for diagnosis of metabolic syndrome. Each parameter has to be interpreted based on the proposed cut-offs, making the diagnosis slightly complex and error-prone. This package is developed by incorporating the modified ATP-III guidelines, and it will aid in the easy and quick diagnosis of metabolic syndrome in busy healthcare settings and also for research purposes. The modified ATP-III-AHA-NHLBI criteria for the diagnosis is described by Grundy et al ., (2005) <doi:10.1161/CIRCULATIONAHA.105.169404>.

Usage

```
MetabolicSyndrome(x)
```

Arguments

x a data frame with column names as exactly specified.

Value

Yes or No

Examples

```
MetabolicSyndrome(x)
```

x

An example data frame

Description

A data frame with exact column names as specified.

Usage

x

Format

A data frame with parameters needed to diagnose metabolic syndrome.

Gender Gender in Male or Female

WC Waist Circumference in cm

TG Triglycerides in mg/dL

HDL High Density Lipoprotein Cholesterol in mg/dL

SBP Systolic BP in mm Hg

DBP Diastolic BP in mm Hg

FPG Fasting plasma glucose in mg/dL

Index

* **datasets**

x, [3](#)

MetabolicSyndrome, [2](#)

x, [3](#)