

Package ‘PowerSDI’

January 20, 2025

Type Package

Title Calculate Standardised Drought Indices Using NASA POWER Data

Version 1.0.0

Description A set of functions designed to calculate the standardised precipitation and standardised precipitation evapotranspiration indices using NASA POWER data as described in Blain et al. (2023) [<doi:10.2139/ssrn.4442843>](https://doi.org/10.2139/ssrn.4442843). These indices are calculated using a reference data source. The functions verify if the indices' estimates meet the assumption of normality and how well NASA POWER estimates represent real-world data. Indices are calculated in a routine mode. Potential evapotranspiration amounts and the difference between rainfall and potential evapotranspiration are also calculated. The functions adopt a basic time scale that splits each month into four periods. Days 1 to 7, days 8 to 14, days 15 to 21, and days 22 to 28, 29, 30, or 31, where 'TS=4' corresponds to a 1-month length moving window (calculated 4 times per month) and 'TS=48' corresponds to a 12-month length moving window (calculated 4 times per month).

License MIT + file LICENSE

Encoding UTF-8

LazyData true

RoxygenNote 7.2.3

Depends R (>= 3.1.0)

Imports graphics, lmom, lubridate, nasapower, stats

URL <https://github.com/gabrielblain/PowerSDI>

BugReports <https://github.com/gabrielblain/PowerSDI/issues>

Suggests knitr, rmarkdown, testthat (>= 3.0.0), vcr (>= 0.6.0), vdiff (>= 1.0.0)

Config/testthat/edition 3

Config/testthat/parallel true

VignetteBuilder knitr

Language en-GB

NeedsCompilation no

Author Gabriel Constantino Blain [aut, cre]

(<https://orcid.org/0000-0001-8832-7734>),

Graciela R. Sobierajski [aut] (<https://orcid.org/0000-0002-7211-9268>),

Leticia L. Martins [aut] (<https://orcid.org/0000-0002-0299-3005>),

Adam H. Sparks [aut] (<https://orcid.org/0000-0002-0061-8359>)

Maintainer Gabriel Constantino Blain <gabriel.blain@sp.gov.br>

Repository CRAN

Date/Publication 2024-01-15 11:20:02 UTC

Contents

Accuracy	2
DistPar	3
ObsEst	4
OperatSDI	5
plot.PowerSDI.Accuracy	6
PlotData	7
print.PowerSDI.Accuracy	8
Reference	8
refHS	9
refPM	10
ScientSDI	11
Index	14

Accuracy	<i>Verify how well NASA-POWER Data Represent Observed Data</i>
----------	--

Description

Calculates scalar measures of accuracy.

Usage

```
Accuracy(obs_est, conf.int = "Yes", sig.level = 0.95)
```

Arguments

obs_est	A 2-column matrix. The reference or observed and the estimated or predicted data. See ObsEst object as an example.
conf.int	A character variable (Yes or No) defining if the function must calculate confidence intervals. Default is "Yes".
sig.level	A numeric variable (between 0.90 and 0.95) defining the significance level for parameter the confidence intervals. Default is 0.95.

Value

An object of class PowerSDI . Accuracy, a list, which contains:

- Absolute mean error (AME),
- square root of the mean squared error (RMSE),
- Willmott's indices of agreement:
 - original (dorig),
 - modified (dmod) and
 - refined (dref)
- Pearson determination coefficient (R2), and
- if `conf.int = "Yes"`, confidence intervals.

Examples

```
a <- Accuracy(obs_est = ObsEst, conf.int = "No")
a

# A generic plotting method is also supplied
plot(a)
```

DistPar	<i>Parameters for Calculating the SDIs Provided by the ScientSDI Function</i>
---------	---

Description

Contains parameters of the gamma and GEV distributions and the $\Pr(\text{Rain} = 0)$, "probzero.rain".

Usage

```
DistPar
```

Format

A data.frame with 13 variables and 48 rows.

lon longitude in decimal degrees

lat latitude in decimal degrees

quart.month The quartile of each month

alfa.rain Shape parameter of the gamma distribution

beta.rain Scale parameter of the gamma distribution

probzero.rain Probability of rain=0

loc.harg Location parameter of the GEV distribution, PE calculated by HS method

sc.harg Scale parameter of the GEV distribution, PE calculated by HS method
sh.harg Shape parameter of the GEV distribution, PE calculated by HS method
loc.pm Location parameter of the GEV distribution, PE calculated by PM method
sc.pm Scale parameter of the GEV distribution, PE calculated by PM method
sh.pm Shape parameter of the GEV distribution, PE calculated by PM method
TS Time scale at which the SDIs will be calculated

Source

Generated by the `ScientSDI()` function using NASA POWER data.

Examples

```
data(DistPar)
```

ObsEst

Example Data of the Input Required by the Accuracy Function

Description

Contains pairs of reference and estimated data.

Usage

```
ObsEst
```

Format

A `data.frame` with 2 variables and 1434 rows.

PE_obs PE data from a reference weather station

PE_est PE data from the NASA POWER project

Source

Generated by the `PowerSDI` package using data from NASA POWER and Agronomic Institute.

Examples

```
data(ObsEst)
```

OperatSDI

*Calculate Routine NASA-SPI and NASA-SPEI Estimates***Description**

Calculates the SPI (Standardized Precipitation Index) and SPEI (Standardized Precipitation-Evapotranspiration Index) using NASA POWER data.

Usage

```
OperatSDI(
  lon,
  lat,
  start.date,
  end.date,
  PEMethod = "HS",
  distr = "GEV",
  parms,
  TS = 4L
)
```

Arguments

lon	longitude in decimal degrees.
lat	latitude in decimal degrees.
start.date	Date at each time when the calculation must start ("YYYY-MM-DD").
end.date	Date at each time when the calculation must end ("YYYY-MM-DD").
PEMethod	A character variable ("HS" (Hargreaves & Samani) or "PM" (Penman-Monteith) defining the potential evapotranspiration method. Default is "HS".
distr	A character variable ("GEV" or "GLO") defining which distribution is used to calculate the SPEI. Default is "GEV" (generalized extreme value) with "GLO" (generalized logistic distributions) as an option.
parms	Parameters required for calculating the SPI and SPEI. It is provided by the ScientSDI function's DistPar.
TS	Time scale on the "quart.month" basis (integer values between 1 and 96).

Value

A data frame with six columns

- rainfall,
- potential evapotranspiration (PE),
- difference between rainfall and PE (in millimeters),
- the NASA-SPI,
- the NASA-SPEI and
- the SDI categories corresponding to each indices estimates.

Examples

```
# This example is not run as it requires data to be downloaded from an API,  
# which may fail. It also uses data included in this package, "DistPar" for  
# `parms` here  
  
OperatSDI(  
  lon = -47.3,  
  lat = -22.67,  
  start.date = "2023-06-01",  
  end.date = "2023-06-30",  
  PMethod = "HS",  
  distr = "GEV",  
  parms = DistPar,  
  TS = 4)
```

plot.PowerSDI.Accuracy

Plots PowerSDI.Accuracy Objects

Description

Custom plot() method for PowerSDI.Accuracy objects.

Usage

```
## S3 method for class 'PowerSDI.Accuracy'  
plot(x, ...)
```

Arguments

x	a 'PowerSDI.Accuracy' object
...	Other parameters as passed to plot()

Value

No return value, called for side effects. Using this will display a scatter plot of reference ETP data (x-axis) and estimated ETP (y-axis) in the active R session.

PlotData

Plot Rainfall and Potential Evapotranspiration Data

Description

Plots rainfall and potential evapotranspiration, both Penman-Monteith and Hargreaves and Samani, amounts using NASA POWER data.

Usage

```
PlotData(lon, lat, start.date, end.date)
```

Arguments

lon	longitude in decimal degrees: (+) Eastern Hemisphere (-) Western Hemisphere.
lat	latitude in decimal degrees: (+) Northern Hemisphere (-) Southern Hemisphere.
start.date	date at which the indices estimates should start ("YYYY-MM-DD").
end.date	date at which the indices estimates should end ("YYYY-MM-DD").

Value

No return value, called for side effects. Using this will display scatter plots of rainfall and potential evapotranspiration accumulated at the 1-quart.month time scale in the active R session.

Examples

```
# This example requires an Internet connection to fetch data and so is only  
# run in interactive sessions
```

```
PlotData(  
  lon = -47.3,  
  lat = -22.87,  
  start.date = "2021-12-28",  
  end.date = "2022-12-31"  
)
```

```
print.PowerSDI.Accuracy
```

Prints PowerSDI.Accuracy Objects

Description

Custom print() method for PowerSDI.Accuracy objects.

Usage

```
## S3 method for class 'PowerSDI.Accuracy'
print(x, digits = max(3L, getOption("digits") - 3L), ...)
```

Arguments

x	a PowerSDI.Accuracy object
digits	The number of digits to be used after the decimal when displaying accuracy values.
...	ignored

Value

Nothing. Side-effect: pretty prints a PowerSDI.Accuracy object in the R console.

Reference

Calculate the SPI and SPEI Using a Reference Data Source

Description

Calculates the Standardised Precipitation Index (SPI) and Standardised Precipitation-Evapotranspiration Index (SPEI) using a reference data source.

Usage

```
Reference(ref, distr = "GEV", PEMethod = "HS", TS = 4L)
```

Arguments

ref	A data frame with the variables required for calculating the SDIs. See refHS or refPM as examples.
distr	A character variable ("GEV" or "GLO") defining which distribution is used to calculate the SPEI. Default is "GEV".
PEMethod	A character variable ("HS" or "PM") defining the potential evapotranspiration method. Default is "HS".
TS	Time scale on the "quart.month" basis (whole values between 1 and 96). Default is 4.

Value

A data frame with five columns

- rain,
- potential evapotranspiration,
- difference between rainfall and potential evapotranspiration,
- SPI calculated at the time scale selected by the user, and
- SPIE calculated at the time scale selected by the user

Examples

```
Reference(ref = refHS, distr = "GEV", PEMethod = "HS", TS = 4)
```

 refHS

Example of the Input Required by the Reference Function

Description

Contains data for calculating the SPI and SPEI.

Usage

```
refHS
```

Format

A data frame with 10950 rows and 8 variables.

YEAR Year

MM Month

DD Day

tavg Daily average air temperature at 2 metres above the ground (degrees C)

tmax Daily maximum air temperature at 2 metres above the ground (degrees C)

tmin Daily minimum air temperature at 2 metres above the ground (degrees C)

Ra Daily top of the atmosphere radiation (MJ/m²/day)

Rain Daily rainfall amounts (mm)

Source

Agronomic Institute and NASA POWER.

Examples

```
data(refHS)
```

refPM

Example of the Input Required by the Reference Function

Description

Contains data for calculating the SPI and SPEI.

Usage

refPM

Format

A data frame with 10958 rows and 11 variables.

YEAR Year

MM Month

DD Day

tavg Daily average air temperature at 2 metres above the ground (degrees C)

tmax Daily maximum air temperature at 2 metres above the ground (degrees C)

tmin Daily minimum air temperature at 2 metres above the ground (degrees C)

Ra Daily top of the atmosphere radiation (MJ/m²/day)

Rs Daily global horizontal irradiance (MJ/m²/day)

W Daily average wind speed at 2 metres above the ground (m/s)

RH Daily average relative humidity at 2 metres above the ground (in percentage)

Rain Daily rainfall amounts (mm)

Source

Agronomic Institute and NASA POWER.

Examples

data(refPM)

ScientSDI	<i>Estimate parameters of Gamma, Generalized Extreme Value, or Generalized Logistic Distributions</i>
-----------	---

Description

Verifies concepts expected from SDI. The first step of the SPI and SPEI algorithms is to calculate the cumulative probabilities of their input variables (Guttman 1999). Function estimates the parameters of the gamma, generalized extreme value (GEV), or generalized logistic distributions (GLO) through the L-moments method are provided. This function also allows users to remove suspicious values from the data sample.

Usage

```
ScientSDI(
  lon,
  lat,
  start.date,
  end.date,
  distr = "GEV",
  TS = 4L,
  Good = "No",
  sig.level = 0.95,
  RainUplim = NULL,
  RainLowlim = NULL,
  PEUplim = NULL,
  PELowlim = NULL
)
```

Arguments

lon	longitude in decimal degrees: (+) Eastern Hemisphere, (-) Western Hemisphere.
lat	latitude in decimal degrees: (+) Northern hemisphere, (-) Southern Hemisphere.
start.date	date at which the indices estimates should start. Format: "YYYY-MM-DD".
end.date	date at which the indices estimates should end. Format: "YYYY-MM-DD".
distr	A character variable ("GEV" or "GLO") defining the distribution to calculate the SPEI. Default is GEV.
TS	Time scale on the quart.month basis (integer values between 1 and 96). Default is 4.
Good	A character variable ("Yes" or "No") to calculate or not the goodness-of-fit and normality tests. Default is "No".
sig.level	A numeric variable (between 0.90 and 0.95) defining the significance level for parameter Good. Default is "0.95".
RainUplim	Optional. Upper limit in millimetres from which rainfall values larger than it will be removed. Default is NULL.

RainLowlim	Optional. Lower limit in millimetres from which rainfall values smaller than it will be removed. Default is NULL.
PEUplim	Optional. Upper limit in millimetres from which evapotranspiration values larger than it will be removed. Valid for Hargreaves and Samani method Default is NULL.
PELowlim	Optional. Lower limit in millimetres from which evapotranspiration values smaller than it will be removed. Valid for Hargreaves and Samani method Default is NULL.

Value

A list object with data calculated at the time scale selected by the user. If Good = "Yes", this list object includes:

SDI The "NASA-SPI", "NASA-SPEI.HS" and "NASA-SPEI.PM."

DistPar The parameters of the distributions (gamma and GEV) used to calculate the indices.

GoodFit The Lilliefors and Anderson-Darling tests goodness-of-fit tests.

Normality The outcomes of the two normality checking procedures (Wu *et al.*, 2006 and Stagge *et al.*, 2015).

If Good = "No", this list object includes SDI and DistPar.

This function also presents other data (in millimeters) calculated from the NASA POWER project:

- Rainfall amounts (Rain),
- potential evapotranspiration values estimated through the Hargreaves and Samani method (PEHS),
- potential evapotranspiration values estimated through the FAO-56 Penman-Monteith method (PEPM), and
- the difference between rainfall and potential evapotranspiration (PPEHS and PPEPM).

References

Guttman, N.B., 1999. Accepting the standardized precipitation index: a calculation algorithm 1. JAWRA Journal of the American Water Resources Association, 35(2), pp.311-322.

Stagge, J.H., Tallaksen, L.M., Gudmundsson, L., Van Loon, A.F. and Stahl, K., 2015. Candidate distributions for climatological drought indices (SPI and SPEI). International Journal of Climatology, 35(13), pp.4027-4040.

Wu, H., Svoboda, M.D., Hayes, M.J., Wilhite, D.A. and Wen, F., 2006. Appropriate application of the standardized precipitation index in arid locations and dry seasons. International Journal of Climatology: A Journal of the Royal Meteorological Society, 27(1), pp.65-79.

Examples

```
# This example requires an Internet connection to fetch data and takes >5s
# to run, and so is only run in interactive sessions
```

```
ScientSDI(  
  lon = -47.3,  
  lat = -22.87,  
  start.date = "1993-01-01",  
  end.date = "2022-12-31",  
  TS = 1,  
  Good = "no"  
)
```

Index

* datasets

DistPar, [3](#)

ObsEst, [4](#)

refHS, [9](#)

refPM, [10](#)

Accuracy, [2](#)

DistPar, [3](#)

ObsEst, [4](#)

OperatSDI, [5](#)

plot.PowerSDI.Accuracy, [6](#)

PlotData, [7](#)

print.PowerSDI.Accuracy, [8](#)

Reference, [8](#)

refHS, [9](#)

refPM, [10](#)

ScientSDI, [5](#), [11](#)