

# Package ‘llogistic’

October 13, 2022

**Title** The L-Logistic Distribution

**Version** 1.0.3

**Description** Density, distribution function, quantile function and random generation for the L-Logistic distribution with parameters  $m$  and  $\phi$ . The parameter  $m$  is the median of the distribution.

**Imports** stats

**Depends** R ( $\geq 3.3.0$ )

**License** GPL-3

**Encoding** UTF-8

**LazyData** true

**RoxygenNote** 6.1.0

**NeedsCompilation** no

**Author** Rosineide Fernando da Paz [aut, cre],  
Jorge Luís Bazán [ctb]

**Maintainer** Rosineide Fernando da Paz <rfpaz2@gmail.com>

**Repository** CRAN

**Date/Publication** 2019-03-31 20:20:03 UTC

## R topics documented:

llogistic . . . . .	1
<b>Index</b>	<b>4</b>

---

llogistic                      *The L-Logistic Distribution*

---

### Description

Density, distribution function, quantile function and random generation for the L-Logistic distribution with parameters  $m$  and  $\phi$ .

**Usage**

```
dllogistic(x, m, phi, log = FALSE)

pllogistic(q, m, phi, lower.tail = TRUE, log.p = FALSE)

qllogistic(p, m, phi, lower.tail = TRUE, log.p = FALSE)

rlllogistic(n, m, phi)
```

**Arguments**

<code>x, q</code>	vector of quantiles.
<code>m, phi</code>	parameters of the L-Logistic distribution. The parameter <code>m</code> lies in the interval (0,1) and <code>phi</code> is positive.
<code>log, log.p</code>	logical; if TRUE, probabilities <code>p</code> are given as $\log(p)$ .
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P[X \leq x]$ , otherwise, $P[X > x]$ .
<code>p</code>	vector of probabilities.
<code>n</code>	number of observations.

**Details**

The llogistic distribution has density

$$f(x) = \phi(1-m)^{\phi} m^{\phi} x^{\phi} (1-x)^{\phi} (\phi-1) / ((1-m)^{\phi} x^{\phi} + m^{\phi} (1-x)^{\phi})^2,$$

for  $0 < x < 1$ , where `m` is a median of the distribution and `phi` is a shape parameter.

**Value**

`dllogistic(x,m,phi)` gives the density function, `rlllogistic(n,m,phi)` gives `n` random variates and `qllogistic(p,m,phi)` gives the quantile.

**Source**

The L-Logistic distribution was introduced by Tadikamalla and Johnson (1982), which refer to this distribution as Logit-Logistic distribution. Here, we have a new parameterization of the Logit-Logistic with the median as a parameter.

**References**

Paz, R.F., Balakrishnan, N and Bazán, Jorge L. (2016). L-Logistic Distribution: Properties, Inference and an Application to Study Poverty and Inequality in Brazil. São Carlos: Universidade Federal de São Carlos. Technical-Scientific Report No. 261, Theory and Method. Sponsored by the Department of Statistical, <URL:<http://www.pipges.ufscar.br/publicacoes/relatorios-tecnicos/arquivos-1/rt261.pdf>>.

TADIKAMALLA, P. R.; JOHNSON, N. L. (1982). Systems of frequency curves generated by transformations of logistic variables. *Biometrika*, v. 69, n. 2, p. 461.

**Examples**

```
dllogistic(0.3, 0.5, 2)  
pllogistic(0.7, 0.5, 2)  
qllogistic(0.2, 0.5, 2)  
rllogistic(10, 0.5, 2)
```

# Index

`dlogistic(llogistic)`, 1

`llogistic`, 1

`plogistic(llogistic)`, 1

`qllogistic(llogistic)`, 1

`rllogistic(llogistic)`, 1