



df\_sordes

*Example Data Frame for adsorption, desorption, and supply parameter***Description**

User is advised to prepare the data as suggested in the example to fit adsorption or desorption data pertaining to a particular nutrient, e.g., phosphorus (as phosphate), sulphur (as sulphate), micronutrient cations or anions, silicon (as silicate), etc. in soil to the functions namely, Freundlich\_A, Freundlich\_D, Langmuir, and SP. However, to use these functions, researchers should carry out the adsorption-desorption study in the method described as follows: W g (e.g., 2 g) of processed soil samples are taken in a series of 50 mL centrifuge tubes (polypropylene) each containing V mL (e.g., 20 mL) 0.1 M sodium chloride solution with different levels (e.g., 10, 20, 40, 60, ... mg/L) of the nutrient under consideration (e.g., P). The tubes are shaken continuously for a defined period (e.g., 24 h) using a mechanical shaker. Immediately after that, each of them is centrifuged (e.g., at 10000 rpm) for sufficient time (e.g., 10-12 min), and a certain volume (e.g., 15 mL) of clear aliquot is pipetted out and filtered through a Whatman No. 42 filter paper. Then the nutrient content (e.g., P) in the filtrate is estimated by standard procedures. The difference between the quantities of the nutrient in the bathing solution before (initial concentration) and after equilibration (equilibrium or final concentration) is taken as the amount of the nutrient adsorbed by the soil from bathing solution. For desorption study, once supernatant is removed after completion of the adsorption step, a certain volume (e.g., 15 mL) of 0.1 M sodium chloride solution is added in the same sample to reach a final volume (V<sub>final</sub>). For convenience in calculation, V<sub>final</sub> should be equal to V (as taken initially). Generally, soil sample with the highest level of added nutrient from the adsorption study is used in desorption experiment. With the help of mechanical shaker, the soil is re-suspended and equilibrated for 12 h. After shaking, the tubes are centrifuged (e.g., at 10000 rpm) for sufficient time (e.g., 10–12 min), and the same volume (as in the previous step, e.g., 15 mL) of clear supernatant solution is pipetted out for determination of the concentration of the nutrient. This process can be repeated multiple times depending on the objective and convenience of the researcher. The difference between equilibrium concentration at a particular desorption step and equilibrium concentration previous to that desorption step, multiplied by solution:soil ratio (e.g., 20:2) is considered as desorbed amount of the nutrient. The adsorbed amount remaining after each desorption step can be computed by subtracting the desorbed amount at a particular step from the amount of adsorbed nutrient present before that desorption step.

**Usage**

df\_sordes

**Format**

Write the following notations on the spreadsheet: Initial\_conc for Initial concentration (mg/L) of the added element, e.g., phosphorus Equilibrium\_conc for Final or equilibrium concentration (mg/L) after adsorption of the same element Cf\_Des for Final or equilibrium concentration (mg/L) after desorption of the same element

---

 Freundlich\_A

*Freundlich isotherm fitted to adsorption data*


---

**Description**

The linear form of Freundlich adsorption isotherm can be fitted to adsorption data to find out the empirical constants of Freundlich equation (Freundlich, 1926).

**Usage**

```
Freundlich_A(W = W, V = V, Ci = Ci, Cf = Cf, ...)
```

**Arguments**

W	Mass of soil sample (g)
V	Volume of extractant solution (mL)
Ci	Initial concentration (mg/L) of the added element, e.g., phosphorus
Cf	Final or equilibrium concentration (mg/L) after adsorption of the same element
...	Any other argument that can be passed to base plot

**Details**

Freundlich equation/isotherm is used to study the adsorption behavior of any element in soils (Freundlich, 1926). It is of the general form:  $x/m = ac^{1/n}$ , where 'x' is the amount of the adsorbate (e.g., P, Ni, etc.) adsorbed on 'm' amount of the adsorbent (e.g., soil), and 'c' is the equilibrium concentration of the adsorbate. This adsorption model helps to understand the relationship between quantity of any element adsorbed per unit soil weight and their concentration in soil solution. Freundlich equation does not predict or include maximum adsorption capacity of soil, so better suited to dilute solutions (of the element under consideration) in contact with the adsorbent. The 'a' and '1/n' are two empirical-constants, sensitive to the given adsorbent-adsorbate system and temperature. Freundlich 'a' indicates the binding affinity of any element in soil. The '1/n' is the exponent of the equilibrium concentration term in the Freundlich equation, and is always  $< 1$  indicating that the energy of adsorption decreases logarithmically as the fraction of adsorbent-surface covered by the adsorbate increases with increasing equilibrium concentration.

**Value**

a - empirical constant (unitless) 1/n - empirical constant (unitless)

**References**

Freundlich, H., 1926. Colloid and Capillary Chemistry. London: Methuen.

**Examples**

```
with(data = df_sordes, Freundlich_A(W = 2, V = 20, Ci = Initial_conc,
Cf = Equilibrium_conc))
```

---

 Freundlich\_D

*Freundlich isotherm for desorption data in soil*


---

### Description

This function fits Freundlich adsorption isotherm to data pertaining to desorption of an adsorbed nutrient, e.g., phosphorus (as phosphate), sulphur (as sulphate), micronutrient cations or anions, silicon (as silicate), etc. in soil.

### Usage

```
Freundlich_D(W = W, V = V, Ci = Ci, Cf = Cf, Cf_Des = Cf_Des, Vres= Vres,
Vfinal = Vfinal, onebyn_ads = onebyn_ads, ...)
```

### Arguments

W	Mass of soil sample (g)
V	Volume of extractant solution (mL)
Ci	Initial concentration (mg/L) of the added element, e.g., phosphorus
Cf	Final or equilibrium concentration (mg/L) after adsorption of the same element
Cf_Des	Final or equilibrium concentration (mg/L) after each desorption step of the same element
Vres	Volume of carried over solution from previous adsorption or desorption step (mL)
Vfinal	Final volume of the solution for each desorption step (mL)
onebyn_ads	Values of '1/n' obtained from Freundlich_A
...	Any other argument that can be passed to base plot

### Value

a - empirical constant (unitless) 1/n - empirical constant (unitless)

### References

Barman, M., Datta, S.P., Rattan, R.K., Meena, M.C., 2013. Sorption and desorption of nickel in soils in relation to its availability to plants. *Agrochimica LVII* (3), 235–249.

### Examples

```
with(data = df_sordes, Freundlich_D(W = 2, V = 20, Ci = Initial_conc, Cf = Equilibrium_conc,
Cf_Des = Cf_Des, Vres= 5, Vfinal = 20, onebyn_ads = 0.2056263))
```

Langmuir

*Langmuir isotherm fitted to adsorption data***Description**

The linear form of Langmuir adsorption isotherm can be fitted to adsorption data to find out adsorption maxima, affinity coefficient, and maximum buffering capacity (Langmuir, 1918).

**Usage**

Langmuir(W = W, V = V, Ci = Ci, Cf = Cf, ...)

**Arguments**

W	Mass of soil sample (g)
V	Volume of extractant solution (mL)
Ci	Initial concentration (mg/L) of the added element, e.g., phosphorus
Cf	Final or equilibrium concentration (mg/L) after adsorption of the same element
...	Any other argument that can be passed to base plot

**Value**

b - Maximum monolayer adsorption or adsorption maxima (mg/kg) k - Constant related to binding energy or the affinity coefficient for the nutrient, e.g., P (L/mg) MBC - Maximum buffering capacity of soil for the nutrient under consideration (L/kg)

**References**

Havlin, J.L., Tisdale, S.L., Nelson, W.L., Beaton, J.D., 2016. Phosphorus. In: Soil fertility and fertilizers. Pearson Education India, pp.160–198. Holford, I.C.R., Mattingly, G.E.G., 1976. Phosphate adsorption and plant availability of phosphate. *Plant and Soil* 44, 377–389. <https://doi.org/10.1007/BF00015889>  
 Langmuir, I., 1918. The adsorption of gases on plane surfaces of glass, mica, and platinum. *Journal of American Chemical Society* 40, 1361–1403. <https://doi.org/10.1021/ja02242a004> Shirvani, M., Shariatmadari, H., Kalbasi, M., 2005. Phosphorus buffering capacity indices as related to soil properties and plant uptake. *Journal of Plant Nutrition*, 28, 537–550. <https://doi.org/10.1081/PLN-200049235>

**Examples**

```
with(data = df_sordes, Langmuir(W = 2, V = 20, Ci = Initial_conc, Cf = Equilibrium_conc))
```

---

SP *Supply parameter of phosphorus in soil*

---

### Description

This function generates the supply parameter (SP) of phosphorus in soil as described by Khasawneh and Copeland (1973).

### Usage

`SP(W = W, V = V, Ci = Ci, Cf = Cf, ...)`

### Arguments

W	Mass of soil sample (g)
V	Volume of extractant solution (mL)
Ci	Initial concentration (mg/L) of the added element, e.g., phosphorus
Cf	Final or equilibrium concentration (mg/L) after adsorption of the same element
...	Any other argument that can be passed to base plot

### Value

SP - Supply parameter  $(\text{mg}^{0.5}/(\text{kg L})^{0.25})$

### References

Khasawneh, F.E., 1971. Solution ion activity and plant growth. *Soil Science Society of America Proceedings* 35, 426–436. Khasawneh, F.E., Copeland, J.P., 1973. Cotton root growth and uptake of nutrients: relation of phosphorus uptake to quantity, intensity, and buffering capacity. *Soil Science Society of America Proceedings* 37, 250–254.

### Examples

```
with(data = df_sordes, SP(W = 2, V = 20, Ci = Initial_conc, Cf = Equilibrium_conc,
col = rgb(red = 1, green = 0, blue = 0, alpha = 0.5), pch = 16, cex = 1))
```

# Index

\* **datasets**

df\_sordes, [2](#)

df\_sordes, [2](#)

Freundlich\_A, [3](#)

Freundlich\_D, [4](#)

Langmuir, [5](#)

SP, [6](#)