

Package ‘SpatialPOP’

January 20, 2025

Type Package

Title Generation of Spatial Data with Spatially Varying Model
Parameter

Version 0.1.0

Author Nobin Chandra Paul

Maintainer Nobin Chandra Paul<ncp375@gmail.com>

Depends R(>= 2.10)

Suggests knitr, rmarkdown, testthat (>= 3.0.0)

VignetteBuilder knitr

Description A spatial population can be generated based on spatially varying regression model under the assumption that observations are collected from a uniform two-dimensional grid consist of $(m * m)$ lattice points with unit distance between any two neighbouring points. For method details see Chao, Liu., Chuanhua, Wei. and Yunnan, Su. (2018).<[DOI:10.1080/10485252.2018.1499907](https://doi.org/10.1080/10485252.2018.1499907)>. This spatially generated data can be used to test different issues related to the statistical analysis of spatial data. This generated spatial data can be utilized in geographically weighted regression analysis for studying the spatially varying relationships among the variables.

License GPL (>= 2.0)

Encoding UTF-8

RoxygenNote 7.1.2

Imports base,MASS,stats, qpdf, numbers

NeedsCompilation no

Repository CRAN

Date/Publication 2022-04-01 07:50:02 UTC

Contents

spatialPOP	2
spatial_grid	3

Index	4
--------------	----------

spatialPOP	<i>Generation of Spatial Data with Spatial Coordinates and Spatially Varying Model Parameters</i>
------------	---

Description

Generation of Spatial Data with Spatial Coordinates and Spatially Varying Model Parameters

Usage

```
spatialPOP(N, m, xlat, ylong)
```

Arguments

N	integer; population size i.e. $N = (m * m)$
m	integer
xlat	numeric vector
ylong	numeric vector

Value

returns a dataframe of spatially generated population consist of simulated response variable (i.e. Y) along with their spatial coordinates, spatially varying model parameters and one explanatory variable (i.e. X)

References

1. Leung, Y., Mei, C. L. and Zhang, W. X. (2000). Statistical tests for spatial non-stationarity based on the geographically weighted regression model. *Environment and Planning A*, 32(1), 9-32.
2. Chao, Liu., Chuanhua, Wei. and Yunan, Su. (2018). Geographically weighted regression model-assisted estimation in survey sampling. *Journal of Nonparametric Statistics*. <DOI:10.1080/10485252.2018.1499907>.

Examples

```
lattice_points<-spatial_grid(c(1:10),c(1:10))  
spatial_data<-spatialPOP(100,10,c(1:10),c(1:10))
```

<code>spatial_grid</code>	<i>a uniform two dimensional grid of lattice points</i>
---------------------------	---

Description

a uniform two dimensional grid of lattice points

Usage

```
spatial_grid(lat, long)
```

Arguments

<code>lat</code>	numeric vector
<code>long</code>	numeric vector

Value

returns a dataframe of lattice points

Examples

```
spatial_grid(c(1:10),c(1:10))
```

Index

spatial_grid, 3
spatialPOP, 2