

# Package ‘LIStest’

January 20, 2025

**Type** Package

**Title** Tests of independence based on the Longest Increasing Subsequence

**Version** 2.1

**Date** 2014-03-12

**Author** J. E. Garcia and V. A. Gonzalez-Lopez

**Maintainer** J. E. Garcia <jg@ime.unicamp.br>

**Depends** R (>= 2.10)

**Description** Tests for independence between X and Y computed from a paired sample  $(x_1, y_1), \dots, (x_n, y_n)$  of  $(X, Y)$ , using one of the following statistics (a) the Longest Increasing Subsequence ( $L_n$ ), (b)  $JL_n$ , a Jackknife version of  $L_n$  or (c)  $JLM_n$ , a Jackknife version of the longest monotonic subsequence. This family of tests can be applied under the assumption of continuity of X and Y.

**License** GPL-2

**LazyLoad** yes

**LazyData** yes

**NeedsCompilation** no

**Repository** CRAN

**Date/Publication** 2014-03-12 23:17:20

## Contents

LIStest-package . . . . .	2
JLMn . . . . .	2
JLn . . . . .	3
lis . . . . .	5
lis.test . . . . .	6
Ln . . . . .	7
TJLMN . . . . .	8
TJLN . . . . .	9
TLN . . . . .	9

<b>Index</b>	<b>10</b>
--------------	-----------

---

LISestest-package

*Tests of independence based on the Longest Increasing Subsequence*

---

### Description

Tests for independence between  $X$  and  $Y$  computed from a paired sample  $(x_1, y_1), \dots, (x_n, y_n)$  of  $(X, Y)$ , using one of the following statistics (a) the Longest Increasing Subsequence (Ln), (b) JLn, a Jackknife version of Ln or (c) JLMn, a Jackknife version of the longest monotonic subsequence. This family of tests can be applied under the assumption of continuity of  $X$  and  $Y$ .

### Details

Package: LISestest  
Type: Package  
Version: 2.1  
Date: 2014-03-12  
License: GPL-2

### Author(s)

J. E. Garcia and V. A. Gonzalez-Lopez Maintainer: J. E. Garcia <jg@ime.unicamp.br>

### References

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

---

JLMn

*JLMn statistic, to test independence*

---

### Description

It compute the JLMn-statistic, from a bivariate sample of continuous random variables  $X$  and  $Y$ .

### Usage

JLMn(x, y)

### Arguments

x, y                    numeric vectors of data values. x and y must have the same length.

**Details**

See subsection 3.3-Main reference. For sample sizes less than 20, the correction introduced in subsection 3.2 from main reference, with  $c = 0.4$  was avoided.

**Value**

The value of the JLMn-statistic.

**Author(s)**

J. E. Garcia, V. A. Gonzalez-Lopez

**References**

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

**Examples**

```
# mixture of two bivariate normal, one with correlation 0.9 and
# the other with correlation -0.9
#
N <-100
ro<- 0.90
Z1<-rnorm(N)
Z2<-rnorm(N)
X2<-X1<-Z1
I<-(1:floor(N*0.5))
I2<-((floor(N*0.5)+1):N)
X1[I]<-Z1[I]
X2[I]<-(Z1[I]*ro+Z2[I]*sqrt(1-ro*ro))
X1[I2]<-Z1[I2]
X2[I2]<-(Z1[I2]*(-ro)+Z2[I2]*sqrt(1-ro*ro))
plot(X1,X2)

#calculate the statistic
a<-JLMn(X1,X2)
a
```

---

JLn

*JLn statistic, to test independence*

---

**Description**

It compute the JLn-statistic, from a bivariate sample of continuous random variables X and Y.

**Usage**

JLn(x, y)

**Arguments**

x, y                    numeric vectors of data values. x and y must have the same length.

**Details**

See subsection 3.2.-Main reference. For sample sizes less than 20, the correction introduced in subsection 3.2 from main reference, with  $c = 0.4$  was avoided.

**Value**

The value of the JLn-statistic.

**Author(s)**

J. E. Garcia and V. A. Gonzalez-Lopez

**References**

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

**Examples**

```
## mixture of two bivariate normal, one with correlation 0.9 and
## the other with correlation -0.9
#
N <- 100
ro <- 0.90
Z1 <- rnorm(N)
Z2 <- rnorm(N)
X2 <- X1 <- Z1
I <- (1:floor(N*0.5))
I2 <- ((floor(N*0.5)+1):N)
X1[I] <- Z1[I]
X2[I] <- (Z1[I]*ro+Z2[I]*sqrt(1-ro*ro))
X1[I2] <- Z1[I2]
X2[I2] <- (Z1[I2]*(-ro)+Z2[I2]*sqrt(1-ro*ro))
plot(X1,X2)

# calculate the statistic
a <- JLn(X1,X2)
a
```

---

`lis`*Longest increasing subsequence for a univariate sample*

---

**Description**

It compute the size of the longest increasing subsequence from a sample of a (continuous) random variable.

**Usage**`lis(x)`**Arguments**

`x` numeric vector of data values.

**Details**

See example 2.1-Main reference.

**Value**

Integer, the size of the longest increasing subsequence.

**Author(s)**

J. E. Garcia and V. A. Gonzalez-Lopez

**References**

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

**Examples**

```
#see Example 2.1 (reference)
a<-lis(c(3,6,1,7,4,2,5,8))
a
```

lis.test

*Test for independence between paired samples***Description**

Test for independence between X and Y computed from a paired sample  $(x_1, y_1), \dots, (x_n, y_n)$  of  $(X, Y)$ , using one of the following statistics (a) the Longest Increasing Subsequence (Ln), (b) JLn, a Jackknife version of Ln or (c) JLMn, a Jackknife version of the longest monotonic subsequence. This family of tests can be applied under the assumption of continuity of X and Y.

**Usage**

```
lis.test(x, y, alternative = c("two.sided", "less", "greater"),
method = c("JLMn", "Ln", "JLn"))
```

**Arguments**

x, y	numeric vectors of data values. x and y must have the same length.
alternative	indicates the alternative hypothesis and must be one of "two.sided"(default), "greater" or "less".
method	a character string indicating which statistics is to be used for the test. One of "Ln", "JLn", or "JLMn"(default).

**Details**

For sample sizes less than 20, the correction introduced in subsection 3.2 from main reference, with  $c = 0.4$  was avoided.

**Value**

sample.estimate	the value of the statistic.
p.value	the p-value for the test.
alternative	a character string describing the alternative hypothesis.
method	a character string indicating what type of Lis-test was performed.

**Author(s)**

J. E. Garcia and V. A. Gonzalez-Lopez

**References**

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

**Examples**

```

# Example 1
# mixture of two bivariate normal, one with correlation 0.9
# and the other with correlation -0.9

N <- 100
ro <- 0.90
Z1 <- rnorm(N)
Z2 <- rnorm(N)
X2 <- X1 <- Z1
I <- (1:floor(N*0.5))
I2 <- ((floor(N*0.5)+1):N)
X1[I] <- Z1[I]
X2[I] <- (Z1[I]*ro+Z2[I]*sqrt(1-ro*ro))
X1[I2] <- Z1[I2]
X2[I2] <- (Z1[I2]*(-ro)+Z2[I2]*sqrt(1-ro*ro))
plot(X1,X2)
# calculate the p.value using the default settings (method="JLMn"
# and alternative="two.sided")
lis.test(X1,X2)
# calculate the p.value using method="JLn" and
# alternative="two.sided".
lis.test(X1,X2,method="JLn")
#
# Example 2: see subsection 4.3.2-Application 2 from main reference.
# (It requires the package VGAM)
#
#require(VGAM)
#plot(coalminers$BW, coalminers$nBW)
#lis.test(coalminers$BW, coalminers$nBW,
#alternative = "greater", method = "Ln")
#lis.test(coalminers$BW, coalminers$nBW,
#alternative = "greater", method = "JLn")
#

```

---

Ln

*Ln (Longest Increasing Subsequence) statistic, to test independence*


---

**Description**

It compute the Ln-statistic, from a bivariate sample of continuous random variables X and Y.

**Usage**

Ln(x, y)

**Arguments**

x, y                    numeric vectors of data values. x and y must have the same length.

**Details**

See Section 2.-Main reference.

**Value**

The value of the Ln-statistic.

**Author(s)**

J. E. Garcia and V. A. Gonzalez-Lopez

**References**

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

**Examples**

```
## mixture of two bivariate normal, one with correlation
## 0.9 and the other with correlation -0.9
#
N <-100
ro<- 0.90
Z1<-rnorm(N)
Z2<-rnorm(N)
X2<-X1<-Z1
I<-(1:floor(N*0.5))
I2<-((floor(N*0.5)+1):N)
X1[I]<-Z1[I]
X2[I]<-(Z1[I]*ro+Z2[I]*sqrt(1-ro*ro))
X1[I2]<-Z1[I2]
X2[I2]<-(Z1[I2]*(-ro)+Z2[I2]*sqrt(1-ro*ro))
plot(X1,X2)

# calculate the statistic
a<-Ln(X1,X2)
a
```

---

TJLMN

---

*Simulated values for the JLMn statistic*


---

**Description**

Simulated values for the JLMn statistic under the hypothesis of independence

**Format**

The format is: List of 200 tables



**References**

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

---

TJLN

*Simulated values for the JLn statistic*

---

**Description**

Simulated values for the JLn statistic under the hypothesis of independence.

**Format**

The format is: List of 200 tables

**References**

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

---

TLN

*Simulated values for the Ln statistic*

---

**Description**

Simulated values for the Ln statistic under the hypothesis of independence

**Format**

The format is: List of 200 tables

**References**

J. E. Garcia, V. A. Gonzalez-Lopez, Independence tests for continuous random variables based on the longest increasing subsequence, Journal of Multivariate Analysis (2014), <http://dx.doi.org/10.1016/j.jmva.2014.02.010>

# Index

## \* ~copula

- JLMn, 2
- JLn, 3
- lis, 5
- lis.test, 6
- Ln, 7

## \* ~longest increasing subsequence

- JLMn, 2
- JLn, 3
- lis, 5
- lis.test, 6
- Ln, 7

## \* datasets

- TJLMN, 8
- TJLN, 9
- TLN, 9

JLMn, 2

JLn, 3

lis, 5

lis.test, 6

LIStest (LISest-package), 2

LISest-package, 2

Ln, 7

TJLMN, 8

TJLN, 9

TLN, 9