

Package ‘WHORiskCalculator’

April 7, 2026

Title WHO Cardiovascular Disease Risk Calculator

Version 1.0.0

Description Implements the 2019 World Health Organization (WHO) cardiovascular disease (CVD) risk prediction models, as described in Kaptoge et al. (2019) [doi:10.1016/S2214-109X\(19\)30318-3](https://doi.org/10.1016/S2214-109X(19)30318-3). Provides two validated models for estimating 10-year risk of fatal and non-fatal cardiovascular events (myocardial infarction and stroke): a laboratory-based model using age, sex, systolic blood pressure, total cholesterol, smoking status, and diabetes history; and a non-laboratory-based model substituting body mass index (BMI) for cholesterol and diabetes, suitable for resource-limited settings. Risk estimates are recalibrated to 21 Global Burden of Disease regions using region-specific incidence rates and risk factor distributions derived from the Emerging Risk Factors Collaboration. Functions are fully vectorized for efficient batch calculations and support automatic country-to-region mapping via ISO 3166-1 alpha-3 country codes.

License MIT + file LICENSE

Encoding UTF-8

RoxygenNote 7.3.3

Suggests devtools, renv, spelling

Language en-US

Config/testthat/edition 3

NeedsCompilation no

Author WHO CVD Risk Chart Working Group [aut],
Andrea Pedot [cre, ctb]

Maintainer Andrea Pedot <andrea.pedot@unimi.it>

Repository CRAN

Date/Publication 2026-04-07 07:30:09 UTC

Contents

calculate_who_risk	2
------------------------------	---

calculate_who_risk_nonlab	4
country_to_region	6
get_country_codes	6
get_regions	7

Index	8
--------------	----------

calculate_who_risk	<i>Calculate WHO 10-Year CVD Risk (Laboratory-Based Model)</i>
--------------------	--

Description

Calculates the 10-year risk of cardiovascular disease (fatal and non-fatal myocardial infarction and stroke) using the WHO laboratory-based model. This model requires total cholesterol measurement.

Usage

```
calculate_who_risk(
  age,
  sex,
  sbp,
  cholesterol,
  smoking,
  diabetes,
  region = NULL,
  country = NULL
)
```

Arguments

age	Numeric vector. Age in years (valid range: 40-80).
sex	Character vector. Sex: "male" or "female".
sbp	Numeric vector. Systolic blood pressure in mmHg.
cholesterol	Numeric vector. Total cholesterol in mmol/L. To convert from mg/dL to mmol/L, divide by 38.67.
smoking	Logical vector. Current smoking status (TRUE = current smoker).
diabetes	Logical vector. History of diabetes (TRUE = diabetic).
region	Character vector. WHO/GBD region name. One of the 21 global regions (see get_regions for valid values). If NULL and country is provided, region will be determined automatically.
country	Character vector. ISO 3166-1 alpha-3 country code (e.g., "USA", "GBR", "IND"). Used to determine region if region is NULL. See get_country_codes for valid values.

Details

The model uses Cox proportional hazards regression with the following predictors:

- Age (with interactions for all other predictors)
- Total cholesterol
- Systolic blood pressure
- Diabetes status
- Current smoking status

Risk is calculated separately for myocardial infarction/CHD death and stroke, then combined assuming independence of the two outcomes:

$$P_{CVD} = 1 - (1 - P_{MI}) \times (1 - P_{stroke})$$

The model is recalibrated to region-specific incidence rates and mean risk factor levels from the Global Burden of Disease study.

Value

Numeric vector of 10-year CVD risk as a proportion (0-1). Multiply by 100 to get percentage.

Centering Values

Variables are centered at:

- Age: 60 years
- Systolic blood pressure: 120 mmHg
- Total cholesterol: 6 mmol/L

See Also

[calculate_who_risk_nonlab](#) for the non-laboratory-based model, [get_regions](#) for available regions, [get_country_codes](#) for country code lookup.

Examples

```
# Single calculation
calculate_who_risk(
  age = 55,
  sex = "male",
  sbp = 140,
  cholesterol = 5.5,
  smoking = TRUE,
  diabetes = FALSE,
  region = "western_europe"
)

# Using country code instead of region
calculate_who_risk(
```

```

age = 60,
sex = "female",
sbp = 130,
cholesterol = 6.0,
smoking = FALSE,
diabetes = TRUE,
country = "ITA"
)

# Vectorized calculation
calculate_who_risk(
  age = c(45, 55, 65),
  sex = c("male", "female", "male"),
  sbp = c(120, 140, 160),
  cholesterol = c(5.0, 6.0, 7.0),
  smoking = c(FALSE, TRUE, FALSE),
  diabetes = c(FALSE, FALSE, TRUE),
  country = c("USA", "GBR", "IND")
)

```

```
calculate_who_risk_nonlab
```

Calculate WHO 10-Year CVD Risk (Non-Laboratory-Based Model)

Description

Calculates the 10-year risk of cardiovascular disease using the WHO non-laboratory-based model. This model uses BMI instead of cholesterol and does not require diabetes status, making it suitable for resource-limited settings where laboratory testing may not be available.

Usage

```

calculate_who_risk_nonlab(
  age,
  sex,
  sbp,
  bmi,
  smoking,
  region = NULL,
  country = NULL
)

```

Arguments

age	Numeric vector. Age in years (valid range: 40-80).
sex	Character vector. Sex: "male" or "female".
sbp	Numeric vector. Systolic blood pressure in mmHg.

bmi	Numeric vector. Body mass index in kg/m ² .
smoking	Logical vector. Current smoking status (TRUE = current smoker).
region	Character vector. WHO/GBD region name. One of the 21 global regions (see get_regions for valid values). If NULL and country is provided, region will be determined automatically.
country	Character vector. ISO 3166-1 alpha-3 country code.

Details

This model is designed for use in resource-limited settings where laboratory measurements (cholesterol, glucose) may not be readily available. It uses BMI as a proxy for metabolic risk.

Important limitations:

- The non-laboratory model tends to underestimate risk in people with diabetes because diabetes is a strong predictor not included in this model.
- For individuals identified as high risk (>10%) by this model, laboratory-based assessment is recommended.

Value

Numeric vector of 10-year CVD risk as a proportion (0-1).

Centering Values

Variables are centered at:

- Age: 60 years
- Systolic blood pressure: 120 mmHg
- BMI: 25 kg/m²

See Also

[calculate_who_risk](#) for the laboratory-based model.

Examples

```
# Single calculation
calculate_who_risk_nonlab(
  age = 55,
  sex = "male",
  sbp = 140,
  bmi = 28,
  smoking = TRUE,
  region = "south_asia"
)

# Using country code
calculate_who_risk_nonlab(
  age = 60,
```

```
sex = "female",
sbp = 135,
bmi = 26,
smoking = FALSE,
country = "KEN"
)
```

country_to_region *Map Country Code to Region*

Description

Utility function to map ISO 3166-1 alpha-3 country codes to WHO/GBD regions.

Usage

```
country_to_region(country)
```

Arguments

country Character vector of ISO 3166-1 alpha-3 country codes.

Value

Character vector of region names.

Examples

```
country_to_region(c("USA", "DEU", "JPN"))
```

get_country_codes *Get Country Code to Region Mapping*

Description

Returns a named vector mapping ISO 3166-1 alpha-3 country codes to their corresponding WHO/GBD regions.

Usage

```
get_country_codes(country = NULL)
```

Arguments

country Optional character vector of country codes to look up. If NULL (default), returns the complete mapping.

Value

If country is NULL, returns a named character vector where names are country codes and values are region names. If country is provided, returns the corresponding region(s).

Examples

```
# Get all mappings
head(get_country_codes())

# Look up specific countries
get_country_codes(c("USA", "GBR", "IND"))
```

get_regions

Get Available Region Names

Description

Returns a character vector of valid WHO/GBD region names that can be used with the risk calculation functions.

Usage

```
get_regions()
```

Value

Character vector of 21 GBD region names.

Examples

```
get_regions()
```

Index

`calculate_who_risk`, [2](#), [5](#)
`calculate_who_risk_nonlab`, [3](#), [4](#)
`country_to_region`, [6](#)

`get_country_codes`, [2](#), [3](#), [6](#)
`get_regions`, [2](#), [3](#), [5](#), [7](#)